

Modifiable lifestyle factors for primary prevention of chronic kidney disease: a systematic review and meta-analysis

Kelly, Jaimon T; Su, Guobin ; Zhang, La; Qin, Xindong ; Marshall, Skye; Gonzalez-Ortiz, Ailema; Clase, Catherine M; Campbell, Katrina Louise; Xu, Hong; Carrero, Juan-Jesus

Published in:
Journal of the American Society of Nephrology : JASN

DOI:
[10.1681/ASN.2020030384](https://doi.org/10.1681/ASN.2020030384)

Licence:
Other

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Kelly, J. T., Su, G., Zhang, L., Qin, X., Marshall, S., Gonzalez-Ortiz, A., Clase, C. M., Campbell, K. L., Xu, H., & Carrero, J-J. (2021). Modifiable lifestyle factors for primary prevention of chronic kidney disease: a systematic review and meta-analysis. *Journal of the American Society of Nephrology : JASN*, 32(1), 239-253.
<https://doi.org/10.1681/ASN.2020030384>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Modifiable lifestyle factors for primary prevention of chronic kidney disease: a systematic review and meta-analysis

Jaimon T Kelly,¹ Guobin Su,^{2,3} La Zhang,³ Xindong Qin,³ Skye Marshall,⁴ Ailema González-Ortiz,^{2,5} Catherine M Clase⁶, Katrina L Campbell,¹ Hong Xu,^{2,7} Juan-Jesus Carrero²

Affiliations

1. Menzies Health Institute Queensland, Griffith University, Gold Coast, Queensland, Australia
2. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
3. Department of Nephrology, Guangdong Provincial Hospital of Chinese Medicine, The Second Affiliated Hospital, Guangzhou University of Chinese Medicine, Guangzhou city, Guangdong Province, China
4. Bond University Nutrition & Dietetics Research Group, Faculty of Health Science and Medicine, Bond University, Gold Coast, Queensland, Australia; and Nutrition Research Australia, Sydney, New South Wales, Australia.
5. Nephrology and Mineral Metabolism Department, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico
6. Department of Medicine, Department of Health Research Methods, Evidence & Impact, McMaster University, Hamilton, Ontario, Canada
7. Division of Clinical Geriatrics, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Stockholm, Sweden

Keywords: diet, exercise, smoking, alcohol, lifestyle, chronic kidney disease, albuminuria, primary prevention, systematic review

Funding: This study received no specific funding.

Word Count (body): 4,240 (excluding title page, abstract, tables, and references)

Word Count (abstract): 249

Corresponding Authors:

Jaimon T. Kelly, PhD

Menzies Health Institute Queensland, Faculty of Medicine, Griffith University, Australia

jaimon.kelly@griffith.edu.au

and

Juan J. Carrero, PhD

Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Sweden

Juan.jesus.carrero@ki.se

Abstract

Background

Despite the growth of chronic kidney disease (CKD), no evidence-based lifestyle recommendations for primary CKD prevention exist.

Objective

To evaluate the consistency of evidence associating modifiable lifestyle factors and the incidence of CKD.

Methods

MEDLINE, Embase, CINAHL, and references from eligible studies were searched from database inception through June 2019. Cohort studies of adults without CKD at baseline, reporting lifestyle exposures (diet, physical activity, alcohol consumption and tobacco smoking). The primary outcome was incident CKD (estimated glomerular filtration rate [eGFR] <60mL/min/1.73m²). Secondary outcomes were other surrogate measures of CKD, including kidney replacement therapy, GFR decline, and albuminuria.

Results

We identified 104 studies of 2,755,719 participants with generally a low risk of bias. Higher potassium (odds ratio 0.78 [95% CI: 0.65-0.94]; I²=48%, 7 studies) and vegetable intake (0.79 [0.70-0.90]; I²=57%, 5 studies) significantly decreased the odds of CKD, whereas higher salt intake (1.21 [1.06-1.38]; I²=59%, 5 studies) significantly increased the odds. Being physically active had lower odds of CKD (0.82 [0.69-0.98]; I²=83%, 9 studies) compared with being sedentary. Current and former smokers (risk ratio 1.20 [1.12-1.28]; I²=80%, 12 studies) had significantly increased risk of CKD compared with those who never smoked. Compared with no consumption, moderate consumption of alcohol was associated with reduced risk of CKD

(0.86 [0.79-0.93]; $I^2=40\%$, 14 studies). Results for the secondary outcomes were consistent with the primary finding.

Conclusion

This study identified modifiable lifestyle hazards that consistently predict the incidence of CKD in the community and may inform both public health recommendations and clinical practice.

Introduction

Chronic kidney disease (CKD) affects 10% of the world's population¹ and ranks in the top 10 non-communicable diseases contributing to disease and disability.² Its incidence is increasing worldwide,¹ and mortality owing to CKD rose between 2005 and 2017 from 0.9 million to 1.2 million deaths annually². CKD imposes a significant economic burden on patients and society; many developed countries spend between 2-3% of their annual healthcare expenditure on kidney replacement therapies (KRT, that is, chronic dialysis or kidney transplantation) for 0.2 percent of the total population.³

Because prevention is more effective than cure, avoiding exposures to hazards that may cause CKD in the community is a key priority. This is particularly important because the management of patients with established CKD may require dietary adaptations that diametrically differ from those that are needed for primary prevention.⁴ However, no evidence-based lifestyle recommendations for the primary prevention of CKD are apparent. Instead, current advice extrapolates from recommendations and inferences drawn from literature concerning cardiovascular disease, hypertension, and diabetes.⁵⁻⁷ Furthermore, the World Health Organization discusses CKD as a complication of cardiovascular disease and diabetes, which further complicates effective public health messages.⁸ To address this critical knowledge gap, we hereby evaluated the consistency of evidence associating modifiable lifestyle factors (specifically diet, physical activity, alcohol consumption and tobacco smoking) and the incidence of CKD.

Methods

This systematic review and meta-analysis was conducted according to Meta-analysis of Observational Studies in Epidemiology (MOOSE) checklist for observational studies⁹ and was prospectively registered in PROSPERO (CRD42019137328).

Data Sources and Searches

We searched MEDLINE, Embase and CINAHL (database inception through June 2019) without date or language restriction (Supplemental Table 1). We used EndNote to de-duplicate and manage citations. For publications in languages other than English, we used translations by native speakers and where not possible, Google Translate. Author pairs completed the title and abstract screening, full-text screening, and manually searched the reference lists of eligible studies and reviews to identify other potentially relevant studies by snowballing. We included a third investigator (J.K. or J.J.C.) and used consensus to resolve disagreements.

Study Selection

Inclusion criteria were: adult participants without established CKD (i.e., glomerular filtration rate [GFR] $>60\text{mL/min}^2$) at baseline,¹⁰ including those that reported people without CKD as a subgroup of a larger study; cohort studies, prospective or retrospective; and outcome data (primary or secondary) reported based on levels of exposure to a lifestyle factor, including diet (foods or nutrients), physical activity, alcohol consumption and tobacco smoking. We excluded studies of participants with albuminuria at baseline, whenever this information was available. We did not consider diet pattern exposures in our review, as these have recently been systematically evaluated.^{11,12}

The primary outcome was incident CKD, defined as the development of GFR $<60\text{ mL/min/1.73 m}^2$ during follow-up.¹⁰ Secondary outcomes were other surrogate measures of CKD: KRT, GFR decline, or albuminuria.

Data Extraction and Quality Assessment

One author from each team extracted the data from the included studies using standardized data extraction forms created in Microsoft Excel, which was checked by a second author before analysis. Supplemental Table 2 shows the data extracted for analysis. Where we identified more

than one publication for a single cohort study reporting on the same lifestyle factor, we used the data set from the published study contributing the highest number of participants. Within each meta-analysis, we conducted a sensitivity analysis examining the effects of substituting these alternative sources of data on the same cohort. Where questions arose concerning a study's eligibility, missing data, or further information on results, we contacted corresponding authors to request this information. We used author responses to make decision on a study's eligibility and analysis approach.

We assessed risk of bias for each included study, using the Newcastle-Ottawa Quality Assessment Scale,¹³ which assesses the methodological quality of the studies across three domains: selection (or representativeness) of cohorts, comparability (of cohorts due to design or analysis) and outcomes (assessment and follow up) (Supplemental Table 2). We applied The Grading of Recommendations Assessment, Development and Evaluation methodology (GRADE) criteria to rate the quality of evidence (low, moderate or high) for each outcome. In analyses that included five or more studies, when the primary outcome showed low or moderate heterogeneity, we constructed funnel plots to examine for effects that may represent publication, selection or reporting bias.

Data Synthesis and Analysis

We used RevMan 5.3 and Stata version 16.0 (StataCorp, College Station, TX) to meta-analyze data when a particular lifestyle factor was reported for the same outcome from 3 separate cohorts; otherwise associations were tabulated and reported using frequencies. Given the observational design and highly varied sample sizes of the primary studies, we used the random-effects model (DerSimonian & Laird) for all meta-analyses. Robustness of the association estimate was evaluated by comparing if any statistically significant association identified through random-effect models became non-significant under a fixed-effects model.

The overall association estimates for all binary outcomes are expressed as odds ratios (OR) or

relative risks (RR), depending on which ratio was predominantly used in the original data, together with 95% confidence intervals (CI). For studies reporting changes in mean values at the end of their observation follow-up periods, we extracted and tabulated the end of the study values with their associated variance. If the association compared a lifestyle factor with a reference exposure that was not homogenous with the other included lifestyle exposures (e.g., high adherence vs low adherence) then the ratio methods were inverted and then combined into the meta-analysis. We used Chi-square testing at an alpha of 0.05, and with the I^2 statistic, rated low (<25%), moderate (25-75%) or high (>75%) to assess for heterogeneity. Supplemental Table 2 gives details of planned subgroup and sensitivity analyses to explore reasons for heterogeneity. In addition to these linear analyses for alcohol consumption, we also conducted a dose-response random-effects meta-regression analysis to account for potential changes in the risk estimate at different dosages of alcohol consumption (methods reported in Supplemental Table 2).

Results

The electronic search retrieved 45,111 citations of which, a total of 104 studies encompassing 2,755,719 participants met the inclusion criteria (Supplemental Figure 1). The characteristics of included studies are reported in Supplemental Table 3. Lifestyle exposures and associations to kidney outcomes varied across the studies, with n=56 reporting diet exposures, n=18 reporting physical activity, n=27 reporting alcohol consumption and n=30 reporting tobacco smoking. The outcomes evaluated included incident CKD in (n=51 studies, KRT in 17 studies, GFR decline in 32 studies, and albuminuria in 20 studies).

Lifestyle hazards and incident chronic kidney disease

The association between lifestyle factors and incident CKD was reported in 51 studies of 1,221,018¹⁴⁻⁷². The main results of the meta-analysis are summarized in Supplemental Figure 2.

Diet. The association between 57 different dietary factors with incident CKD was described in 31 studies of 176,625 participants.^{14,20,24,26,27,30,38,39,41,46,47,49,51,57,60,61,66,69,70} Meta-analysis was possible for 9 dietary factors (Table 1), of which higher vegetable intake (OR 0.79 [95% CI: 0.70, 0.90]; $I^2=57\%$; Evidence quality: Low) and higher potassium intake (OR 0.78 [95% CI: 0.65, 0.94]; $I^2=48\%$; Evidence quality: Low) were consistently associated with lower odds of CKD. Higher sodium intake was associated with higher odds of CKD (OR 1.21 [95% CI: 1.06, 1.38]; $I^2=59\%$; Evidence quality: Moderate) (Table 1, Figure 1). Meta-analysis of studies evaluating carbohydrate intake, fish, fruit, phosphorus, protein and sugar-sweetened beverages consumption showed no clear association (Table 1).

We were unable to pool data on 32 additional dietary factors reported across 21 studies (Supplemental Table 4).^{15,16,18,19,23,25,27,30-32,37,45-47,57,59-62,71,73} We identified 19 dietary factors for which one or both available studies showed an association with in the direction indicating decreased risk: cereal fiber, coffee, dairy, fiber, folate, legumes, magnesium, nitrate, nuts, nuts & legumes, plant protein, omega-3, DHA, EPA, vitamin B12, vitamin C, vitamin D, vitamin E, and zinc. We identified 2 dietary factors for which one or both available studies indicating a harmful relationship: a higher sodium-potassium ratio and red & processed meat consumption (Supplemental Table 4).

Physical activity. The association of different levels of physical activity with incident CKD was described in 10 studies of 78,301 participants (Table 1; Figure 2).^{28,34,42,44,53,54,58,59,63,72} Meta-analysis of 9 studies showed lower odds of CKD in people who were more physically active compared with those who were less physically active (OR 0.82 [95% CI: 0.69, 0.98]; $I^2=83\%$; Evidence quality: Very low).

Alcohol intake. The association between alcohol intake and incident CKD was described in 14 studies of 211,072 participants.^{22,23,28,40,43,50,52,55,63-65,67,68,72} Compared with lower intakes, meta-analysis showed that higher consumption of alcohol (RR 0.87 [95% CI: 0.79, 0.95]; $I^2=43\%$; Evidence quality: Low) and moderate alcohol consumption (RR 0.86 [95% CI: 0.79, 0.93]; $I^2=50\%$; Evidence quality: Moderate) were associated with lower risk of CKD (Table 1, Figure 3). In a dose–response meta-regression analysis, we modelled the relationship between incident CKD and alcohol intake using restricted cubic splines (Supplemental Table 5 and Supplemental Figure 3) and found that higher alcohol intake was associated with lower risk of incident CKD. This association was generally observed throughout the whole range of alcohol consumption considered with a significant effect size (p values for nonlinearity = 0.03). There was no apparent association when people who had never consumed alcohol were compared with those who had consumed it in the past (former intake) (Table 1).

Smoking. The association between smoking and incident CKD was described in 12 studies of 985,086 participants.^{28,29,35,43,50,52,53,59,63,65,68,74} Compared with people who never smoked, current and former smokers (ever smokers) showed higher odds of CKD (OR 1.18 [95% CI: 1.10, 1.27]; $I^2=81\%$; Evidence quality: Very low) (Table 1, Figure 3). Compared with people who had never smoked, former smokers had increased odds of CKD (OR 1.09 [95% CI: 1.01, 1.17]; $I^2=90\%$; Evidence quality: Very low) in 6 studies of 944,931 participants (Table 1). There was no consistent association with CKD in 2 studies comparing current with former smokers (Supplemental Table 4).

Lifestyle risk factors and secondary outcomes (KRT, GFR decline, and albuminuria):

The consistency of associations between the lifestyle factors identified in our primary analysis and other markers of kidney damage is shown in Figure 4.

The association between lifestyle factors and KRT was described in 17 studies of 990,723 participants;^{33,35,37,75-88} lifestyle factors and GFR decline in 32 studies of 108,936

participants,^{14,20,23,34,46,49,73,86,89-105} and lifestyle factors and albuminuria in 20 studies of 512,403 participants.^{23,39,41,88,95,100,101,105-116} Table 2 and the ‘Supplementary Material. Data synthesis for secondary outcomes’ show the details. The risk of KRT was higher among current and former smokers (RR 1.59 [95% CI: 1.30, 1.94]; $I^2=68\%$; Evidence quality: Moderate) compared with never-smokers (Supplemental Figure 4). The risk of GFR decline was lower in persons with higher potassium intake (RR 0.49 [95% CI: 0.31, 0.79]; $I^2=90\%$; Evidence quality: Very low) and physically active (OR 0.76 [95% CI: 0.59, 0.97]; 75%; Evidence quality: Very low) (Supplemental Figure 5). The risk of albuminuria was lower among physically active persons (OR 0.88 [95% CI: 0.81, 0.96]; $I^2=0\%$; Evidence quality: Low) and higher among tobacco smokers (OR 1.67 [95% CI: 1.23, 2.26]; $I^2=88\%$; Evidence quality: Very low) (Supplemental Figure 6).

Other lifestyle hazards did not have sufficient studies to allow meta-analysis, but their associations suggested they may be potentially protective, potentially harmful or having no association to secondary study outcomes based on the direction of the majority ($\geq 50\%$) of studies. Results are shown in Supplemental Figure 7 and fully detailed in Supplemental Tables 3 & 4. In brief, coffee consumption was associated with lower risk of incident CKD and KRT in 2/2 and 1/2 of studies, respectively. Dairy intake was associated with lower risk of incident CKD and albuminuria in 2/4 and 1/2 of studies, respectively. Red and processed meats was associated with lower risk of incident CKD, KRT and albuminuria in 1/2, 1/1 and 1/1 of studies, respectively.

Subgroup and sensitivity analyses

Subgroup analyses did not show consistent relationships with duration of exposure or geographic region. Replacing individual datasets (substituting alternative reports from the same cohort) did not meaningfully change results (see ‘Supplementary Material. Data synthesis for secondary outcomes’). We also conducted a *post-hoc* sensitivity analysis on the way that

the frequency of alcohol consumption was reported: higher ‘daily’ consumption of alcohol had no significant association with incident CKD (7 studies; RR 0.98 [95% CI 0.82, 1.18]; $I^2 = 43\%$), whereas higher weekly alcohol consumption was associated with a lower risk of incident CKD (6 studies; RR 0.82 [95% CI 0.75, 0.90]; $I^2 = 32\%$) (Supplemental Table 6). Subgroup analysis comparing studies with participants baseline GFR ≥ 89.9 -70 mL/min/1.73 to GFR ≥ 90 mL/min/1.73 showed significant associations between study exposures (except for physical activity) and the risk of incident CKD in studies with participants baseline GFR ≥ 89.9 -70 mL/min/1.73. The association of vegetable intake and incident CKD held robust in both baseline GFR subgroups. Whereas in studies reporting participant baseline GFR ≥ 90 mL/min/1.73 (which were 37 out of 104; 12 studies able to be meta-analyzed), these associations did not attain statistical significance. However, the direction and magnitude of observed risks in both strata were very similar (Supplemental Table 6).

Quality assessment

The certainty (GRADE) of the evidence for each outcome meta-analyzed is detailed in Supplemental Table 8. The overall risk of bias across the studies was low (Supplemental Figure 8 & 9). Eighty-eight percent of studies had a low risk of bias for sample, whereas 10% were rated high due to being conducted in populations with established diseases (such as type 2 diabetes or hypertension) which are known to increase the risk of incident CKD. With respect to follow-up, 82% of studies had low risk of bias, bias was judged unclear in 13% of studies that reported less than 4 years follow-up; 3 studies (5%) had a high risk of bias due to very short follow-up periods of less than 2 years. The risk of bias in exposure assessment was rated high 92% of studies lifestyle behaviors were assessed by self-report; objective measures were used in 8 studies (8%). Outcomes were captured using appropriate tools in 88% of studies, and high is 6% of studies. Potential analysis bias was low in 73% of studies, where 23% were unclear, because data were not analyzed using ratio statistics to allow statistically data pooling.

Discussion

This study evaluated the association between modifiable lifestyle factors and incident CKD in the community. In predominately low-to-very-low-certainty evidence, we found higher intake of potassium and vegetables, lower intake of sodium, physical activity, moderate alcohol consumption, and avoidance of tobacco smoking to be consistently associated with lower risk of CKD. We identified consistency with these main results when we examined their association with other surrogates of kidney damage: KRT, GFR decline and albuminuria.

Higher vegetable intake is associated with higher potassium intake: both dietary factors were consistently associated with fewer kidney outcomes. Low vegetable consumption is a strong predictor of mortality in the general population.^{117,118} The mechanism is likely, at least in part, through the intermediary diseases that are themselves risks for CKD: cardiovascular disease, hypertension, diabetes, obesity, and metabolic syndrome.¹¹⁹ Whether there is additionally a more direct relationship between vegetable or potassium intake and kidney health requires further study.

In our analysis, higher potassium intake was associated with reduced incident CKD and GFR decline, adding to the growing evidence-base for the protective association of potassium intake in other chronic conditions. In the Prospective Urban Rural Epidemiology (PURE) study, higher potassium excretion was associated with a lower risk of death and cardiovascular events across more than 18 countries worldwide.¹²⁰ Higher intakes of potassium have also been demonstrated to lower the risk of stroke¹²¹ and lower blood pressure in adults with and without hypertension.^{122,123} In patients in ONTARGET, a high vascular risk cohort, most of whom did not have CKD at baseline, potassium intake was associated with a less rapid GFR decline or need for KRT.¹²⁴ In ONTARGET, the relationship between potassium intake and the outcome did not appear to differ by baseline level of albuminuria, but there was a statistically significant interaction suggesting that the protective association with potassium might not be observed in

patients with GFR <45 mL/min (advanced CKD). In these patients, potassium, and with it, fruits and vegetables are often restricted. When discussing dietary advice, it is critical to distinguish whether we are concerned with prevention of incident CKD, prevention of progression (and at what level of GFR), or management of hyperkalemia and management of the risk of hyperkalemia.¹²⁵

Similarly, higher sodium intake showed a consistent association with increased risk of incident CKD, KRT and GFR decline. This supports the growing evidence from clinical trials, demonstrating the effect of sodium intake on blood pressure, proteinuria and extra-cellular volume in CKD.¹²⁶ Despite existing controversy surrounding reverse causality at extremely low sodium intakes,¹²⁷ our review supports the message that higher sodium intakes are detrimental and public health efforts should be prioritized to reduce population-wide sodium consumption. More work is needed on the optimal level of sodium restriction; our analysis focused on high vs low intake, which in the original studies, the highest exposures of salt defining high intake varied, ranging from ≥ 9.88 to 16.27g/day (sodium ≥ 172 to 283mmol/day).

A healthy diet pattern, characterized by higher intakes of fruit, vegetables, low-fat dairy, fiber and wholegrain and lower in sodium, red meat and sugar has been associated with a 30% reduced odds of kidney damage.^{11,12} Many of the individual diet factors we identified are characteristic of a higher diet quality and overall healthier diet pattern. For example, taking all the data into account, diet factors possibly protective against incident CKD in our current study included fruit, dairy, vegetables, fiber, legumes, nuts, potassium, and unsaturated fatty acids. In contrast, diet factors possibly harmful to incident CKD in our study included excessive carbohydrate intake, high energy intake, red and processed meats, saturated fat, sodium, sodium-potassium ratio and sugar-sweetened beverages. The protective factors, characteristics of a healthy diet, are also strongly associated with reduced risk of overweight and obesity,¹²⁸ which is linked with increased risk of incident CKD.¹²⁹ These associations fundamentally align

with guidelines for general healthy eating and cardiovascular disease prevention, which is helpful and convenient in guiding public health policy for CKD prevention that aligns with prevention of other chronic diseases.¹³⁰

We found that higher levels of physical activity were consistently associated with a lower risk of all study outcomes evaluated. This aligns with the majority of existing studies, which suggest that physical activity is associated with reduced risk of kidney damage, through attenuating cardiovascular disease risk,^{131,132} mortality¹³³ and rapid declines in kidney function.¹³⁴ Whether a relationship independent of cardiovascular and vascular damage truly exists is not known based on current data. Possible mechanisms include reduced blood pressure, improved glycemic control, angiogenesis and vascular regeneration by the up-regulation of endothelial nitric oxide production and other antioxidant enzymes.¹³⁵ While the measurement of physical activity was heterogeneous across the included studies, the most common categorization of higher levels of activity was defined as at least 30 minutes a day.^{44,59} These findings suggest that public health messages should emulate those of primary prevention of cardiovascular disease, to achieve and maintain a moderate level of physical activity of at least 30 minutes a day and avoiding extended periods of being sedentary.^{136,137}

We found an inverse association between alcohol consumption and incident CKD, KRT and GFR decline. While the relationship between alcohol consumption and kidney function decline has not been consistent in the current body of literature, many clinical studies have indeed shown that moderate alcohol consumption is associated with lower occurrence of CKD.^{40,55,64,68} In contrast, existing literature has shown that chronic alcoholism is associated with CKD,^{56,65} leading to the hypothesis that excessive alcohol consumption directly damages the kidney, independent of liver damage.^{138,139} Similarly, some reports included in our review found that heavy drinking, in contrast with moderate drinking, was associated with increased risk of albuminuria and CKD.^{56,105,140} This distinction, that moderate, but not high, alcohol

consumption is associated with lower prevalence of albuminuria and CKD compared with abstinence or with lower intakes, is consistent with a number of other studies.^{139,141,142} There is also the possibility that unadjusted confounders may be at play, for example, social integration as a product of moderate alcohol consumption and overall well-being, which are good for health.¹⁴³ It would appear that despite minor international variations,¹⁴⁴ and the heterogeneous categorization for the higher alcohol consumption (ranging in grams per day (range 20 to 48g/day), grams per week (210g/week), total drinks per day (range >1 to 4 drinks/day), drinks per week (range 5 to >15 drinks/ week), that moderate alcohol consumption in-line with public health guidelines seems safe and unlikely to cause kidney damage in the general population.

Current and former smokers had higher risk of incident CKD compared with those who never smoked. Tobacco smoking has been previously identified as a risk factor for incident CKD and KRT in the general population.¹⁴⁵ While the mechanism is uncertain, smoking may lead to insulin resistance¹⁴⁶ which might explain the large effect size (67% increase in risk) of smoking on risk of albuminuria observed in our study, consistent with previous observations.¹⁴⁷ The harmful effects of smoking may also be mediated by endothelial cell dysfunction, inducing advanced glycation end products due to glycotoxins present in cigarettes, increased vascular permeability and pathological vascular changes of kidney disease, and insulin resistance in diabetic and non-diabetic subjects.¹⁴⁵ Taken together, these risk factors for kidney function decline clearly support a public health message to avoid tobacco smoking to prevent kidney, as well as cardiovascular disease.

This study has strengths as the first evidence synthesis of lifestyle risk and protective factors and primary prevention of CKD: it is broad in scope and we have taken a rigorous approach to meta-analysis. However, this study has limitations. First, there was scarce evidence for some exposures, particularly for diet factors, precluding meta-analysis; for these factors our output

is simply descriptive. Second, there was variation in the definition of the different exposures across the different lifestyle factors, in the degree of adjustment for potential confounders, and in definition and reporting for study outcomes. For example, there were no consistent adjustments for activity, age, education and sex (as reported in Supplemental Table 3) in all studies included in the analyses. Furthermore, 10 different definitions of incident CKD were used in the included studies, with similar heterogeneity in the definitions of the secondary outcomes). This leads to heterogeneity in the analysis; and precludes clear recommendations in terms of possible targets, goals or thresholds. Third, while we were able to perform meta-analysis to account for the known non-linear associations of alcohol consumption and kidney outcomes, we were unaware of any such assumptions for the other lifestyle factors, and the original studies did not commonly report their associations in this way. The original papers usually made a linear assumption and analyzed data with linear models. It is possible that nonlinear effects or threshold effects may be missed in this way. Fourth, because we evaluated cohort studies and not randomized trials, the findings do not imply causality. However, there were no trials of this nature in our search and it is unlikely that studies of sufficient rigor, size, and duration will be conducted, given the resource intensive nature of such trials. We combined RR and OR; for rare events, these are numerically similar though calculated differently. Fifth, we used the NOS to assess bias rather than the more comprehensive ROBINS-I tool¹⁴⁸ because our resources to complete the study were finite. Finally, in order to conduct meta-analysis, each putative risk and protective factor was considered separately and we do not know whether there are quantitatively important interactions between the different factors (e.g., does the effect of carbohydrate intake differ at different levels of physical activity), as no study has evaluated such hypotheses.

Allowing for the limitations of inferences from observational studies, our work suggests the lifestyle factors we have evaluated are probably those that patients have the most control, they

are not the only ones that public health authorities should target. For example, other potentially modifiable hazards known to increase the risk of kidney damage include environmental pollution,^{149,150} excessive body weight related to nutrition,¹²⁹ inappropriate use of medications (including analgesics, antibiotics, antiretrovirals or proton-pump inhibitors)¹⁵¹ and heavy metal exposure and intake (in-particular cadmium).¹⁵² Finally, society has a further role to play in facilitating the more personal choices, through food labelling, alcohol and tobacco labelling, health policy around exercise and policy around the ease of walking and cycling in the built environment and walkability.

We recognize that these findings are based on non-randomized data with risk of residual and unknown confounding, particularly in the setting of studies with low-to-very-low certainty of the evidence. However, the generalizability and overall importance of these findings should not be undermined, as they are well in-line with public health recommendations for preventing related chronic conditions and general healthy lifestyle principles. Therefore, the results can represent an important indication for public policy, advocacy, and be hypothesis-generating for future randomized trials. It is not trivial to ask patients to change their lifestyle. Diet, in particular, is a highly social and cultural issue; furthermore, many patients are constrained by external factors, including economic factors, in their decisions. For these reasons, we believe that lifestyle intervention studies to address unanswered questions examining the whole range of clinically important outcomes, including the incidence and progressions of CKD, continue to be justified.

Conclusion

Increased vegetable and potassium intake, physical activity and moderate alcohol consumption were consistently associated with reduced risk of CKD. Increased dietary salt intake, and tobacco smoking were consistently associated with increased risk of CKD. In the absence of

trial evidence, these findings can inform both public health recommendations and patient-centered discussions in clinical practice.

ACKNOWLEDGEMENTS

This work is endorsed and performed within the European Renal Nutrition working group, an initiative of and supported by the European Renal Association-European Dialysis and Transplantation Association (ERA-EDTA).

FUNDING

J.J.C is supported by the Swedish Research Council (grant number 2019-01059) and the Swedish Heart and Lung Foundation. G.S is supported by ERA-EDTA Young Fellowship Programme and by the Guangdong provincial hospital of Chinese medicine (No. YN2018QL08). A.G.O. is supported by the Mexican Council of Science and Technology (CVU 373297), School of Medicine, Programa de Doctorado en Ciencias Médicas, Odontológicas y de la Salud. HX is supported by Loo and Hans Osterman's Foundation.

Conflict of interest

J.K. has received consultancy fees from Amgen for travel and lecture presentations, and consultancy fees from HealthCert for topics unrelated to the submitted work. J.J.C reports grant funding from AstraZeneca, ViforPharma and Astellas, consulting for Baxter and AstraZeneca, and speaker fees for Abbott, Nutricia, AstraZeneca and ViforPharma, all outside the submitted work. C.M.C. has received consultation, advisory board membership or research funding from the Ontario Ministry of Health, Sanofi, Johnson & Johnson, Pfizer, Leo Pharma, Astellas, Janssen, Amgen, Boehringer-Ingelheim and Baxter. K.L.C. has received consultation, advisory board membership for research funding from Queensland Health, Nestle Health Sciences and the Dietitians Association of Australia. The rest of co-authors have no conflict of interest to report.

Authors' Contributions

JK, and JC contributed to the study conception. JK, SM, HX, GS, AG, XQ and LZ conducted the literature search, extracted data and appraised risk of bias. JK conducted the data analysis. JK was responsible for writing the first draft of the manuscript. JC reviewed the first draft of the manuscript. KC and CC critically reviewed the manuscript. All authors read and approved the final version of the manuscript.

Supplemental Material

Additional Supplemental Material may be found in the online version of this article at the publisher's web-site:

Supplementary Material. Data synthesis for secondary outcomes

Supplemental Table 1. Search terms used across the electronic databases

Supplemental Table 2. Summary of the methods relating to data extraction, risk of bias, meta-regression and planned subgroup and sensitivity analyses

Supplemental Table 3. Characteristics of the included studies

Supplemental Table 4. Lifestyle hazards and kidney disease outcomes from individual studies which could not be statistically pooled into meta-analysis

Supplement Table 5. Summary of the studies between alcohol intake and incident CKD

Supplemental Table 6: Subgroup analysis for Incident CKD

Supplemental Table 7. Results from sensitivity analysis

Supplemental Table 8: GRADE table summarizing the quality of the evidence for each outcome

Supplemental Figure 1. Study flow diagram

Supplemental Figure 2. Summary of the associations between modifiable lifestyle risk and protective factors and risk of incident CKD,

Supplement Figure 3. Dose–response relationship between alcohol intake (gram per day) and incident CKD estimated with a random-effect meta-regression-restricted cubic spline model

Supplemental Figure 4: Association of Tobacco smoking and ESKD

Supplemental Figure 5: Summary of the associations between modifiable lifestyle risk and protective factors and risk of GFR decline

Supplemental Figure 6: Summary of the associations between modifiable lifestyle risk and protective factors and risk of Albuminuria

Supplemental Figure 7. Consistency of associations in lifestyle factors which could not be statistically pooled across the markers of kidney function decline.

Supplemental Figure 8. Risk of bias across the included studies

Supplemental Figure 9: Individual assessment of risk of bias across the included studies

Supplemental Figure 10: Funnel plots for lifestyle hazards and incident CKD

Supplemental Figure 11: Funnel plots for lifestyle hazards and secondary outcomes

References

1. Bello AK, Levin A, Tonelli M, et al. Assessment of global kidney health care status. *JAMA*. 2017;317(18):1864-1881.
2. Bikbov B, Purcell CA, Levey AS, et al. Global, regional, and national burden of chronic kidney disease, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet*. 2020;395(10225):709-733.
3. Bello A, Levin A, Tonelli M, et al. Global Kidney Health Atlas: A report by the International Society of Nephrology on the current state of organization and structures for kidney care across the globe. *Kidney international*. 2017;7(2):145–153.
4. Kelly JT, Campbell KL, Carrero JJ. Primary Versus Secondary Prevention of Chronic Kidney Disease: The Case of Dietary Protein. *Journal of Renal Nutrition*. 2018;28(4):225-228.
5. Åkesson A, Weismayer C, Newby P, Wolk A. Combined effect of low-risk dietary and lifestyle behaviors in primary prevention of myocardial infarction in women. *Archives of internal medicine*. 2007;167(19):2122-2127.
6. Chiuve SE, Rexrode KM, Spiegelman D, Logroscino G, Manson JE, Rimm EB. Primary prevention of stroke by healthy lifestyle. *Circulation*. 2008;118(9):947.
7. Chiuve SE, Fung TT, Rexrode KM, et al. Adherence to a low-risk, healthy lifestyle and risk of sudden cardiac death among women. *Jama*. 2011;306(1):62-69.
8. World Health Organization. Global status report on noncommunicable diseases 2014. In: World Health Organization; 2014.
9. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. *JAMA*. 2000;283(15):2008-2012.

10. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group: KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney International Supplementation* 2013;3(1):1-163.
11. Bach KE, Kelly JT, Palmer SC, Khalesi S, Strippoli GF, Campbell KL. Healthy dietary patterns and incidence of CKD: a meta-analysis of cohort studies. *Clinical Journal of the American Society of Nephrology*. 2019;14(10):1441-1449.
12. van Westing A, Küpers L, Geleijnse J. Diet and Kidney Function: a Literature Review. *Current Hypertension Reports*. 2020;22(2):14.
13. Wells G, Shea B, O'Connell D, et al. Newcastle-Ottawa quality assessment scale cohort studies. In:2014.
14. Araki S, Haneda M, Koya D, et al. Urinary Potassium Excretion and Renal and Cardiovascular Complications in Patients with Type 2 Diabetes and Normal Renal Function. *Clinical Journal of the American Society of Nephrology*. 2015;10(12):2152-2158.
15. Asghari G, Yuzbashian E, Mirmiran P, Azizi F. The association between Dietary Approaches to Stop Hypertension and incidence of chronic kidney disease in adults: the Tehran Lipid and Glucose Study. *Nephrology Dialysis Transplantation*. 2017;32(Supp 2):ii224-ii230.
16. Asghari G, Yuzbashian E, Shahemi S, Gaeini Z, Mirmiran P, Azizi F. Dietary total antioxidant capacity and incidence of chronic kidney disease in subjects with dysglycemia: Tehran Lipid and Glucose Study. *European journal of nutrition*. 2018;57(7):2377-2385.
17. Baggio B, Budakovic A, Perissinotto E, et al. Atherosclerotic risk factors and renal function in the elderly: the role of hyperfibrinogenaemia and smoking. Results from

- the Italian Longitudinal Study on Ageing (ILSA). *Nephrology Dialysis Transplantation*. 2005;20(1):114-123.
18. Bahadoran Z, Mirmiran P, Ghasemi A, Carlstrom M, Azizi F, Hadaegh F. Association between Dietary Intakes of Nitrate and Nitrite and the Risk of Hypertension and Chronic Kidney Disease: Tehran Lipid and Glucose Study. *Nutrients*. 2016;8(12).
 19. Bahadoran Z, Mirmiran P, Golzarand M, Davudabadi-Farahani R, Azizi F. Dietary Animal-derived L-Arginine Intakes and Risk of Chronic Kidney Disease: a 6-year Follow-up of Tehran Lipid and Glucose Study. *Iranian journal of kidney diseases*. 2017;11(5):352-359.
 20. Bahadoran Z, Mirmiran P, Momenan AA, Azizi F. Allium vegetable intakes and the incidence of cardiovascular disease, hypertension, chronic kidney disease, and type 2 diabetes in adults: a longitudinal follow-up study. *Journal of hypertension*. 2017;35(9):1909-1916.
 21. Bomback AS, Derebail VK, Shoham DA, et al. Sugar-sweetened soda consumption, hyperuricemia, and kidney disease. *Kidney international*. 2010;77(7):609-616.
 22. Buja A, Scafato E, Baggio B, et al. Renal impairment and moderate alcohol consumption in the elderly. Results from the Italian Longitudinal Study on Aging (ILSA). *Public Health Nutrition*. 2011;14(11):1907-1918.
 23. Dunkler D, Dehghan M, Teo KK, et al. Diet and kidney disease in high-risk individuals with type 2 diabetes mellitus. *JAMA internal medicine*. 2013;173(18):1682-1692.
 24. Dunkler D, Kohl M, Teo KK, et al. Dietary risk factors for incidence or progression of chronic kidney disease in individuals with type 2 diabetes in the European Union. *Nephrology Dialysis Transplantation*. 2015;30:iv76-85.

25. Ejtahed HS, Angoorani P, Asghari G, Mirmiran P, Azizi F. Dietary Advanced Glycation End Products and Risk of Chronic Kidney Disease. *Journal of Renal Nutrition*. 2016;26(5):308-314.
26. Farhadnejad H, Asghari G, Emamat H, Mirmiran P, Azizi F. Low-Carbohydrate High-Protein Diet is Associated With Increased Risk of Incident Chronic Kidney Diseases Among Tehranian Adults. *Journal of Renal Nutrition*. 2018;29(4):343-349.
27. Farhadnejad H, Asghari G, Mirmiran P, Yuzbashian E, Azizi F. Micronutrient Intakes and Incidence of Chronic Kidney Disease in Adults: Tehran Lipid and Glucose Study. *Nutrients*. 2016;8(4):217.
28. Foster MC, Hwang S-J, Massaro JM, Jacques PF, Fox CS, Chu AY. Lifestyle factors and indices of kidney function in the Framingham Heart Study. *American journal of nephrology*. 2015;41(4-5):267-274.
29. Fox CS, Larson MG, Leip EP, Culleton B, Wilson PW, Levy D. Predictors of new-onset kidney disease in a community-based population. *JAMA*. 2004;291(7):844-850.
30. Gopinath B, Harris DC, Flood VM, Burlutsky G, Brand-Miller J, Mitchell P. Carbohydrate nutrition is associated with the 5-year incidence of chronic kidney disease. *Journal of Nutrition*. 2011;141(3):433-439.
31. Gopinath B, Harris DC, Flood VM, Burlutsky G, Mitchell P. Associations between dairy food consumption and chronic kidney disease in older adults. *Scientific reports*. 2016;6:39532.
32. Haring B, Selvin E, Liang M, et al. Dietary Protein Sources and Risk for Incident Chronic Kidney Disease: Results From the Atherosclerosis Risk in Communities (ARIC) Study. *Journal of Renal Nutrition*. 2017;27(4):233-242.
33. Haroun MK, Jaar BG, Hoffman SC, Comstock GW, Klag MJ, Coresh J. Risk factors for chronic kidney disease: a prospective study of 23,534 men and women in

- Washington County, Maryland. *Journal of the American Society of Nephrology*. 2003;14(11):2934-2941.
34. Hawkins M, Newman AB, Madero M, et al. The associations of physical activity and television watching with change in kidney function in older adults. *Journal of Physical Activity & Health*. 2015;12(4):561.
 35. Hippisley-Cox J, Coupland C. Predicting the risk of Chronic Kidney Disease in Men and Women in England and Wales: prospective derivation and external validation of the QKidney® Scores. *BMC Family Practice*. 2010;11(1):49.
 36. Hu EA, Lazo M, Rosenberg SD, et al. Alcohol consumption and incident kidney disease: results from the atherosclerosis risk in communities study. *Journal of Renal Nutrition*. 2020;30(1):22-30.
 37. Hu EA, Selvin E, Grams ME, Steffen LM, Coresh J, Rebholz CM. Coffee consumption and incident kidney disease: results from the Atherosclerosis Risk in Communities (ARIC) Study. *American Journal of Kidney Diseases*. 2018;72(2):214-222.
 38. Jhee JH, Kee YK, Park JT, et al. A Diet Rich in Vegetables and Fruit and Incident CKD: A Community-Based Prospective Cohort Study. *American Journal of Kidney Diseases*. 2019;74(4):491-500.
 39. Kieneker LM, Bakker SJ, de Boer RA, Navis GJ, Gansevoort RT, Joosten MM. Low potassium excretion but not high sodium excretion is associated with increased risk of developing chronic kidney disease. *Kidney international*. 2016;90(4):888-896.
 40. Koning SH, Gansevoort RT, Mukamal KJ, Rimm EB, Bakker SJ, Joosten MM. Alcohol consumption is inversely associated with the risk of developing chronic kidney disease. *Kidney international*. 2015;87(5):1009-1016.

41. Lee CC, Howard BV, Mete M, Wang H, Jolly S, Adler AI. Association Between Fish Consumption and Nephropathy in American Indians-The Strong Heart Study. *Journal of Renal Nutrition*. 2012;22(2):221-227.
42. Lin H-C, Peng C-H, Chiou J-Y, Huang C-N. Physical activity is associated with decreased incidence of chronic kidney disease in type 2 diabetes patients: A retrospective cohort study in Taiwan. *Primary Care Diabetes*. 2014;8(4):315-321.
43. Michishita R, Matsuda T, Kawakami S, et al. The association between changes in lifestyle behaviors and the incidence of chronic kidney disease (CKD) in middle-aged and older men. *Journal of Epidemiology*. 2017;27(8):389-397.
44. Michishita R, Matsuda T, Kawakami S, et al. The joint impact of habitual exercise and glycemic control on the incidence of chronic kidney disease (CKD) in middle-aged and older males. *Environmental Health and Preventive Medicine*. 2017;22(1):76.
45. Mirmiran P, Bahadoran Z, Golzarand M, Asghari G, Azizi F. Consumption of nitrate containing vegetables and the risk of chronic kidney disease: Tehran Lipid and Glucose Study. *Renal failure*. 2016;38(6):937-944.
46. Mirmiran P, Nazeri P, Bahadoran Z, Khalili-Moghadam S, Azizi F. Dietary Sodium to Potassium Ratio and the Incidence of Chronic Kidney Disease in Adults: A Longitudinal Follow-Up Study. *Preventive nutrition and food science*. 2018;23(2):87-93.
47. Mirmiran P, Yuzbashian E, Asghari G, Sarverzadeh S, Azizi F. Dietary fibre intake in relation to the risk of incident chronic kidney disease. *British Journal of Nutrition*. 2018;119(5):479-485.
48. Miyatake N, Moriyasu H, Sakano N, Tada S, Suzue T, Hirao T. Influence of cigarette smoking on estimated glomerular filtration rate (eGFR) in Japanese male workers. *Acta Medica Okayama*. 2010;64(6):385-390.

49. Mun KH, Yu GI, Choi BY, Kim MK, Shin MH, Shin DH. Association of dietary potassium intake with the development of chronic kidney disease and renal function in patients with mildly decreased kidney function: The Korean multi-rural communities cohort study. *Medical Science Monitor*. 2019;25:1061-1070.
50. Nakanishi N, Fukui M, Tanaka M, et al. Low urine pH is a predictor of chronic kidney disease. *Kidney and Blood Pressure Research*. 2012;35(2):77-81.
51. Nam KH, An SY, Joo YS, et al. Carbohydrate-Rich Diet Is Associated with Increased Risk of Incident Chronic Kidney Disease in Non-Diabetic Subjects. *Journal of clinical medicine*. 2019;8(6).
52. Noborisaka Y, Ishizaki M, Yamada Y, et al. The effects of continuing and discontinuing smoking on the development of chronic kidney disease (CKD) in the healthy middle-aged working population in Japan. *Environmental Health and Preventive Medicine*. 2013;18(1):24-32.
53. Obermayr RP, Temml C, Knechtelsdorfer M, et al. Predictors of new-onset decline in kidney function in a general middle-European population. *Nephrology Dialysis Transplantation*. 2008;23(4):1265-1273.
54. Ogunmoroti O, Allen NB, Cushman M, et al. Association between Life's Simple 7 and noncardiovascular disease: the Multi-Ethnic Study of Atherosclerosis. *JAMA*. 2016;5(10):e003954.
55. Okada Y, Uehara S, Shibata M, et al. Habitual Alcohol Intake Modifies Relationship of Uric Acid to Incident Chronic Kidney Disease. *American journal of nephrology*. 2019;50(1):55-62.
56. Pan C-s, Ju TR, Lee CC, et al. Alcohol use disorder tied to development of chronic kidney disease: A nationwide database analysis. *PloS one*. 2018;13(9).

57. Park I, Xun P, Tsinosoi CL, Klemmer P, Liu K, He K. Intakes of long-chain omega-3 polyunsaturated fatty acids and non-fried fish in relation to incidence of chronic kidney disease in young adults: a 25-year follow-up. *European journal of nutrition*. 2019;59(1):399-407.
58. Ramezankhani A, Tohidi M, Azizi F, Hadaegh F. Application of survival tree analysis for exploration of potential interactions between predictors of incident chronic kidney disease: a 15-year follow-up study. *J Transl Med*. 2017;15(1):240-240.
59. Rebholz CM, Anderson CA, Grams ME, et al. Relationship of the AHA impact goals (life's simple 7) with risk of chronic kidney disease: Results from the ARIC cohort study. *Journal of the American Heart Association*. 2016;5(4):e003192.
60. Rebholz CM, Coresh J, Grams ME, et al. Dietary Acid Load and Incident Chronic Kidney Disease: Results from the ARIC Study. *American journal of nephrology*. 2015;42(6):427-435.
61. Rebholz CM, Crews DC, Grams ME, et al. DASH (Dietary Approaches to Stop Hypertension) Diet and Risk of Subsequent Kidney Disease. *American Journal of Kidney Diseases*. 2016;68(6):853-861.
62. Rebholz CM, Young BA, Katz R, et al. Patterns of Beverages Consumed and Risk of Incident Kidney Disease. *Clinical Journal of the American Society of Nephrology*. 2019;14(1):49-56.
63. Ryoo J-H, Choi J-M, Oh C-M, Kim M-G. The association between uric acid and chronic kidney disease in Korean men: a 4-year follow-up study. *Journal of Korean Medical Science*. 2013;28(6):855-860.
64. Schaeffner ES, Kurth T, de Jong PE, Glynn RJ, Buring JE, Gaziano JM. Alcohol consumption and the risk of renal dysfunction in apparently healthy men. *Archives of internal medicine*. 2005;165(9):1048-1053.

65. Shankar A, Klein R, Klein BE. The association among smoking, heavy drinking, and chronic kidney disease. *American Journal of Epidemiology*. 2006;164(3):263-271.
66. Sugiura T, Takase H, Ohte N, Dohi Y. Dietary Salt Intake is a Significant Determinant of Impaired Kidney Function in the General Population. *Kidney and Blood Pressure Research*. 2018;43(4):1245-1254.
67. Weiner DE, Tighiouart H, Elsayed EF, Griffith JL, Salem DN, Levey AS. Uric acid and incident kidney disease in the community. *Journal of the American Society of Nephrology*. 2008;19(6):1204-1211.
68. Yamagata K, Ishida K, Sairenchi T, et al. Risk factors for chronic kidney disease in a community-based population: a 10-year follow-up study. *Kidney international*. 2007;71(2):159-166.
69. Yoon CY, Noh J, Lee J, et al. High and low sodium intakes are associated with incident chronic kidney disease in patients with normal renal function and hypertension. *Kidney international*. 2018;93(4):921-931.
70. Yoon CY, Park JT, Jhee JH, et al. High dietary phosphorus density is a risk factor for incident chronic kidney disease development in diabetic subjects: a community-based prospective cohort study. *American Journal of Clinical Nutrition*. 2017;106(1):311-321.
71. Yuzbashian E, Asghari G, Mirmiran P, Zadeh-Vakili A, Azizi F. Sugar-sweetened beverage consumption and risk of incident chronic kidney disease: Tehran lipid and glucose study. *Nephrology*. 2016;21(7):608-616.
72. Kanda E, Muneyuki T, Suwa K, Nakajima K. Alcohol and exercise affect declining kidney function in healthy males regardless of obesity: A prospective cohort study. *PloS one*. 2015;10(8).

73. Rebholz CM, Tin A, Liu Y, et al. Dietary Magnesium and Kidney Function Decline: The Healthy Aging in Neighborhoods of Diversity across the Life Span Study. *American journal of nephrology*. 2016;44(5):381-387.
74. Tohidi M, Hasheminia M, Mohebi R, et al. Incidence of chronic kidney disease and its risk factors, results of over 10 year follow up in an Iranian cohort. *PloS one*. 2012;7(9).
75. Bash LD, Astor BC, Coresh J. Risk of incident ESRD: a comprehensive look at cardiovascular risk factors and 17 years of follow-up in the Atherosclerosis Risk in Communities (ARIC) Study. *American Journal of Kidney Diseases*. 2010;55(1):31-41.
76. Hallan SI, Orth SR. Smoking is a risk factor in the progression to kidney failure. *Kidney international*. 2011;80(5):516-523.
77. Jafar TH, Jin A, Koh WP, Yuan JM, Chow KY. Physical activity and risk of end-stage kidney disease in the Singapore Chinese Health Study. *Nephrology*. 2015;20(2):61-67.
78. Lew QJ, Jafar TH, Jin A, Yuan JM, Koh WP. Consumption of Coffee but Not of Other Caffeine-Containing Beverages Reduces the Risk of End-Stage Renal Disease in the Singapore Chinese Health Study. *Journal of Nutrition*. 2018;148(8):1315-1322.
79. Lew QJ, Jafar TH, Koh HW, et al. Red Meat Intake and Risk of ESRD. *Journal of the American Society of Nephrology*. 2017;28(1):304-312.
80. Lipworth L, Mumma MT, Cavanaugh KL, et al. Incidence and predictors of end stage renal disease among low-income blacks and whites. *PloS one*. 2012;7(10):e48407.
81. Pscheidt C, Nagel G, Zitt E, Kramar R, Concini H, Lhotta K. Sex-and time-dependent patterns in risk factors of end-stage renal disease: a large Austrian cohort with up to 20 years of follow-up. *PloS one*. 2015;10(8).

82. Rebholz CM, Grams ME, Steffen LM, et al. Diet Soda Consumption and Risk of Incident End Stage Renal Disease. *Clinical Journal of the American Society of Nephrology*. 2017;12(1):79-86.
83. Reynolds K, Gu D, Chen J, et al. Alcohol consumption and the risk of end-stage renal disease among Chinese men. *Kidney international*. 2008;73(7):870-876.
84. Smyth A, Griffin M, Yusuf S, et al. Diet and Major Renal Outcomes: A Prospective Cohort Study. The NIH-AARP Diet and Health Study. *Journal of Renal Nutrition*. 2016;26(5):288-298.
85. Stengel B, Tarver-Carr ME, Powe NR, Eberhardt MS, Brancati FL. Lifestyle factors, obesity and the risk of chronic kidney disease. *Epidemiology*. 2003;14(4):479-487.
86. Van Noordenne ND, Olde Engberink RHG, Van Den Hoek TC, Van Den Born BJH, Vogt L. Associations of 15-year average potassium intake with long term renal outcome in the outpatient clinical setting. *Nephrology Dialysis Transplantation*. 2016;31:i11.
87. Jin A, Koh W-P, Chow KY, Yuan J-M, Jafar TH. Smoking and risk of kidney failure in the Singapore Chinese health study. *PloS one*. 2013;8(5):e62962.
88. Wadén J, Tikkanen HK, Forsblom C, et al. Leisure-time physical activity and development and progression of diabetic nephropathy in type 1 diabetes: the FinnDiane Study. *Diabetologia*. 2015;58(5):929-936.
89. Cirillo M, Cavallo P, Bilancio G, Lombardi C, Vagnarelli OT, Laurenzi M. Low protein intake in the population: Low risk of kidney function decline but high risk of mortality. *Journal of Renal Nutrition*. 2018;28(4):235-244.
90. Deriaz D, Guessous I, Vollenweider P, et al. Estimated 24-h urinary sodium and sodium-to-potassium ratio are predictors of kidney function decline in a population-based study. *Journal of hypertension*. 2019;37(9):1853-1860.

91. Esmeijer K, Geleijnse J, De Fijter J, Kromhout D, Hoogeveen E. High protein intake accelerates kidney function decline in post-myocardial infarction patients-The alpha omega cohort study. *Nephrology Dialysis Transplantation*. 2018;33:i469.
92. Halbesma N, Bakker SJ, Jansen DF, et al. High protein intake associates with cardiovascular events but not with loss of renal function. *Journal of the American Society of Nephrology*. 2009;20(8):1797-1804.
93. Herber-Gast GC, van Essen H, Verschuren WM, et al. Coffee and tea consumption in relation to estimated glomerular filtration rate: results from the population-based longitudinal Doetinchem Cohort Study. *American Journal of Clinical Nutrition*. 2016;103(5):1370-1377.
94. Herber-Gast GCM, Biesbroek S, Verschuren WMM, et al. Association of dietary protein and dairy intakes and change in renal function: Results from the population-based longitudinal Doetinchem cohort study^{1,2}. *American Journal of Clinical Nutrition*. 2016;104(6):1712-1719.
95. Herber-Gast GM, Boersma M, Verschuren WMM, et al. Consumption of whole grains, fruit and vegetables is not associated with indices of renal function in the population-based longitudinal Doetinchem study. *British Journal of Nutrition*. 2017;118(5):375-382.
96. Hirahatake KM, Jacobs DR, Gross MD, et al. The Association of Serum Carotenoids, Tocopherols, and Ascorbic Acid With Rapid Kidney Function Decline: The Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Journal of Renal Nutrition*. 2019;29(1):65-73.
97. Jhee JH, Kee YK, Park S, et al. High-protein diet with renal hyperfiltration is associated with rapid decline rate of renal function: a community-based prospective cohort study. *Nephrology Dialysis Transplantation*. 2019;35(1):98-106.

98. Knight EL, Stampfer MJ, Hankinson SE, Spiegelman D, Curhan GC. The impact of protein intake on renal function decline in women with normal renal function or mild renal insufficiency. *Annals of internal medicine*. 2003;138(6):460-467.
99. Krop JS, Coresh J, Chambless LE, et al. A community-based study of explanatory factors for the excess risk for early renal function decline in blacks vs whites with diabetes: the Atherosclerosis Risk in Communities study. *Archives of Internal Medicine*. 1999;159(15):1777-1783.
100. Lin J, Curhan GC. Associations of sugar and artificially sweetened soda with albuminuria and kidney function decline in women. *Clinical Journal of the American Society of Nephrology*. 2011;6(1):160-166.
101. Lin J, Hu FB, Curhan GC. Associations of diet with albuminuria and kidney function decline. *Clinical Journal of the American Society of Nephrology*. 2010;5(5):836-843.
102. Malhotra R, Lipworth L, Cavanaugh KL, et al. Protein Intake and Long-term Change in Glomerular Filtration Rate in the Jackson Heart Study. *Journal of Renal Nutrition*. 2018;28(4):245-250.
103. Menon V, Katz R, Mukamal K, et al. Alcohol consumption and kidney function decline in the elderly: alcohol and kidney disease. *Nephrology Dialysis Transplantation*. 2010;25(10):3301-3307.
104. Ohta Y, Tsuchihashi T, Kiyohara K, Oniki H. High salt intake promotes a decline in renal function in hypertensive patients: a 10-year observational study. *Hypertension Research*. 2013;36(2):172-176.
105. White SL, Polkinghorne KR, Cass A, Shaw JE, Atkins RC, Chadban SJ. Alcohol consumption and 5-year onset of chronic kidney disease: the AusDiab study. *Nephrology Dialysis Transplantation*. 2009;24(8):2464-2472.

106. Barbato A, D'Elia L, Perna L, et al. Increased Microalbuminuria Risk in Male Cigarette Smokers: Results from the "Olivetti Heart Study" after 8 Years Follow-Up. *Kidney and Blood Pressure Research*. 2019;44(1):33-42.
107. Chang A, Van Horn L, Jacobs DR, Jr., et al. Lifestyle-related factors, obesity, and incident microalbuminuria: the CARDIA (Coronary Artery Risk Development in Young Adults) study. *American Journal of Kidney Diseases*. 2013;62(2):267-275.
108. Forman JP, Scheven L, de Jong PE, Bakker SJ, Curhan GC, Gansevoort RT. Association between sodium intake and change in uric acid, urine albumin excretion, and the risk of developing hypertension. *Circulation*. 2012;125(25):3108-3116.
109. Jee SH, Boulware LE, Guallar E, Suh I, Appel LJ, Miller ER. Direct, progressive association of cardiovascular risk factors with incident proteinuria: results from the Korea Medical Insurance Corporation (KMIC) study. *Archives of Internal Medicine*. 2005;165(19):2299-2304.
110. Kimura Y, Yamamoto R, Shinzawa M, et al. Alcohol consumption and incidence of proteinuria: a retrospective cohort study. *Clinical and Experimental Nephrology*. 2018;22(5):1133-1142.
111. Maeda I, Hayashi T, Sato KK, et al. Cigarette smoking and the association with glomerular hyperfiltration and proteinuria in healthy middle-aged men. *Clinical Journal of the American Society of Nephrology*. 2011;6(10):2462-2469.
112. O'Seaghdha CM, Hwang S-J, Upadhyay A, Meigs JB, Fox CS. Predictors of incident albuminuria in the Framingham Offspring cohort. *American Journal of Kidney Diseases*. 2010;56(5):852-860.
113. Uehara S, Hayashi T, Sato KK, et al. Relationship between alcohol drinking pattern and risk of proteinuria: the Kansai Healthcare Study. *Journal of Epidemiology*. 2016;26(9):464-470.

114. Wakasugi M, Kazama J, Narita I, et al. Association between overall lifestyle changes and the incidence of proteinuria: a population-based, cohort study. *Internal Medicine*. 2017;56(12):1475-1484.
115. Wakasugi M, Kazama JJ, Yamamoto S, Kawamura K, Narita I. A combination of healthy lifestyle factors is associated with a decreased incidence of chronic kidney disease: a population-based cohort study. *Hypertension Research*. 2013;36(4):328-333.
116. Wen J, Hao J, Zhang Y, et al. Fresh fruit consumption and risk of incident albuminuria among rural Chinese adults: A village-based prospective cohort study. *PloS one*. 2018;13(5):e0197917.
117. Wang X, Ouyang Y, Liu J, et al. Fruit and vegetable consumption and mortality from all causes, cardiovascular disease, and cancer: systematic review and dose-response meta-analysis of prospective cohort studies. *BMJ*. 2014;349:g4490.
118. Crowe FL, Roddam AW, Key TJ, et al. Fruit and vegetable intake and mortality from ischaemic heart disease: results from the European Prospective Investigation into Cancer and Nutrition (EPIC)-Heart study. *European heart journal*. 2011;32(10):1235-1243.
119. Chauveau P, Koppe L, Combe C, Lasseur C, Trolonge S, Aparicio M. Vegetarian diets and chronic kidney disease. *Nephrology Dialysis Transplantation*. 2019;34(2):199-207.
120. O'Donnell M, Mente A, Rangarajan S, et al. Joint association of urinary sodium and potassium excretion with cardiovascular events and mortality: prospective cohort study. *BMJ*. 2019;364:l772.

121. Vinceti M, Filippini T, Crippa A, de Sesmaisons A, Wise LA, Orsini N. Meta-analysis of potassium intake and the risk of stroke. *Journal of the American Heart Association*. 2016;5(10):e004210.
122. Whelton PK, He J, Cutler JA, et al. Effects of oral potassium on blood pressure: meta-analysis of randomized controlled clinical trials. *Jama*. 1997;277(20):1624-1632.
123. Geleijnse JM, Kok FJ, Grobbee DE. Blood pressure response to changes in sodium and potassium intake: a metaregression analysis of randomised trials. *Journal of human hypertension*. 2003;17(7):471.
124. Smyth A, Dunkler D, Gao P, et al. The relationship between estimated sodium and potassium excretion and subsequent renal outcomes. *Kidney Int*. 2014;86(6):1205-1212.
125. Clase CM, Carrero J-J, Ellison DH, et al. Potassium homeostasis and management of dyskalemia in kidney diseases: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) Controversies Conference. *Kidney international*. 2020;97(1):42-61.
126. McMahon EJ, Campbell KL, Bauer JD, Mudge DW. Altered dietary salt intake for people with chronic kidney disease. *Cochrane Database of Systematic Reviews*. 2015(2):CD010070.
127. Stolarz-Skrzypek K, Staessen JA. Reducing salt intake for prevention of cardiovascular disease—times are changing. *Adv Chronic Kidney Dis*. 2015;22(2):108-115.
128. Mu M, Xu L-F, Hu D, Wu J, Bai M-J. Dietary Patterns and Overweight/Obesity: A Review Article. *Iran J Public Health*. 2017;46(7):869-876.

129. Garofalo C, Borrelli S, Minutolo R, Chiodini P, De Nicola L, Conte G. A systematic review and meta-analysis suggests obesity predicts onset of chronic kidney disease in the general population. *Kidney international*. 2017;91(5):1224-1235.
130. Mozaffarian D. Dietary and policy priorities for cardiovascular disease, diabetes, and obesity: a comprehensive review. *Circulation*. 2016;133(2):187-225.
131. Kubota Y, Iso H, Yamagishi K, Sawada N, Tsugane S. Daily Total Physical Activity and Incident Cardiovascular Disease in Japanese Men and Women: Japan Public Health Center–Based Prospective Study. *Circulation*. 2017;135(15):1471-1473.
132. Chomistek AK, Cook NR, Rimm EB, Ridker PM, Buring JE, Lee IM. Physical Activity and Incident Cardiovascular Disease in Women: Is the Relation Modified by Level of Global Cardiovascular Risk? *Journal of the American Heart Association*. 2018;7(12):e008234.
133. Lear SA, Hu W, Rangarajan S, et al. The effect of physical activity on mortality and cardiovascular disease in 130 000 people from 17 high-income, middle-income, and low-income countries: the PURE study. *The Lancet*. 2017;390(10113):2643-2654.
134. Robinson-Cohen C, Katz R, Mozaffarian D, et al. Physical activity and rapid decline in kidney function among older adults. *Archives of Internal Medicine*. 2009;169(22):2116-2123.
135. Zelle DM, Klaassen G, van Adrichem E, Bakker SJ, Corpeleijn E, Navis G. Physical inactivity: a risk factor and target for intervention in renal care. *Nature reviews Nephrology*. 2017;13(3):152-168.
136. Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Journal of the American College of Cardiology*. 2019;74(10):e177-e232.

137. Milam RH. Exercise guidelines for chronic kidney disease patients. *Journal of Renal Nutrition*. 2016;26(4):e23-e25.
138. Latchoumycandane C, Nagy LE, McIntyre TM. Chronic ethanol ingestion induces oxidative kidney injury through taurine-inhibitable inflammation. *Free Radical Biology and Medicine*. 2014;69:403-416.
139. Fan Z, Yun J, Yu S, Yang Q, Song L. Alcohol Consumption Can be a "Double-Edged Sword" for Chronic Kidney Disease Patients. *Med Sci Monit*. 2019;25:7059-7072.
140. Lin M, Su Q, Huang H, et al. Alcohol consumption and the risk for renal hyperfiltration in the general Chinese population. *European journal of clinical nutrition*. 2017;71(4):500-505.
141. Roerecke M, Rehm J. Alcohol consumption, drinking patterns, and ischemic heart disease: a narrative review of meta-analyses and a systematic review and meta-analysis of the impact of heavy drinking occasions on risk for moderate drinkers. *BMC medicine*. 2014;12(1):182.
142. Jespersen T, Kruse N, Mehta T, Kuwabara M, Nouredine L, Jalal D. Light wine consumption is associated with a lower odd for cardiovascular disease in chronic kidney disease. *Nutrition, Metabolism and Cardiovascular Diseases*. 2018;28(11):1133-1139.
143. Dunkler D, Kohl M, Heinze G, et al. Modifiable lifestyle and social factors affect chronic kidney disease in high-risk individuals with type 2 diabetes mellitus. *Kidney international*. 2015;87(4):784-791.
144. Baranchuk A, Haseeb S, Alexander B. Alcohol consumption guidelines: international discrepancies and variations. *BMJ*. 2018;361:k1630.

145. Xia J, Wang L, Ma Z, et al. Cigarette smoking and chronic kidney disease in the general population: a systematic review and meta-analysis of prospective cohort studies. *Nephrology Dialysis Transplantation*. 2017;32(3):475-487.
146. Thomas G, Sehgal AR, Kashyap SR, Srinivas TR, Kirwan JP, Navaneethan SD. Metabolic syndrome and kidney disease: a systematic review and meta-analysis. *Clinical Journal of the American Society of Nephrology*. 2011;6(10):2364-2373.
147. Janssen WM, Hillege H, Pinto-Sietsma SJ, Bak AA, De Zeeuw D, de Jong PE. Low levels of urinary albumin excretion are associated with cardiovascular risk factors in the general population. *Clinical Chemistry and Laboratory Medicine*. 2000;38(11):1107-1110.
148. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016;355:i4919.
149. Wei Y, Wang Y, Di Q, et al. Short term exposure to fine particulate matter and hospital admission risks and costs in the Medicare population: time stratified, case crossover study. *BMJ*. 2019;367:l6258.
150. Blum MF, Surapaneni A, Stewart JD, et al. Particulate Matter and Albuminuria, Glomerular Filtration Rate, and Incident CKD. *Clinical Journal of the American Society of Nephrology*. 2020;(in-press).
151. Whittaker CF, Miklich MA, Patel RS, Fink JC. Medication Safety Principles and Practice in CKD. *Clinical Journal of the American Society of Nephrology*. 2018;13(11):1738-1746.
152. Johri N, Jacquillet G, Unwin R. Heavy metal poisoning: the effects of cadmium on the kidney. *Biometals*. 2010;23(5):783-792.

Figure legends

Figure 1. Association of vegetable intake (panel 1A), potassium intake (panel 1B), sodium intake (panel 1C) and incident chronic kidney disease. Note the association estimate for each lifestyle factor is presented on the ratio (OR or RR) which was predominantly used in the included studies.

Figure 2. Association of physical activity level and incident chronic kidney disease.

Figure 3. Association of alcohol consumption (panel 3A), tobacco smoking (panel 3B) and incident chronic kidney disease. Note the association estimate for each lifestyle factor is presented on the ratio (OR or RR) which was predominantly used in the included studies.

Figure 4. Consistency of associations in lifestyle factors across the markers of kidney function decline^a. The 4 studies of sodium intake and GFR decline were not meta-analyzable because 2 of the studies reported means and variance, but not ratio data.

Table 1. Meta-analyses performed in the review showing the association of alcohol consumption, diet, physical activity and smoking exposure to the risk of incident CKD

Exposure	Studies	Participants	Association ratio [95% CI] ^a	Heterogeneity [I ²]	Evidence Quality ^b
Diet Factors					
Fish	3	21,226	OR 0.94 [0.86, 1.02]	I ² =5%	Low
Fruit	4	28,779	OR 0.91 [0.79, 1.06]	I ² =60%	Very low
Vegetables	5	32,054	OR 0.79 [0.70, 0.90]	I ² =57%	Low
Sugar-sweetened beverages	4	22,760	OR 1.45 [0.97, 2.15]	I ² =68%	Very low
Carbohydrates	3	20,238	OR 1.08 [0.85, 1.36]	I ² =63%	Very low
Protein	3	19,835	OR 1.08 [0.91, 1.28]	I ² =47%	Very low
Phosphorus	3	23,466	OR 1.00 [0.75, 1.32]	I ² =72%	Very low
Potassium	7	32,647	OR 0.78 [0.65, 0.94]	I ² =48%	Low
Sodium	6	43,772	RR 1.21 [1.06, 1.38]	I ² =59%	Moderate
Physical Activity					
Physical activity - High vs Low levels	9	70,828	RR 0.82 [0.69, 0.98]	I ² =83%	Very low
Alcohol Consumption					
Alcohol - High vs Low intakes	13	216,291	RR 0.87 [0.79, 0.95]	I ² =43%	Low
Alcohol - Moderate vs Low intakes	7	165,415	RR 0.86 [0.79, 0.93]	I ² =50%	Moderate
Tobacco Smoking					
Tobacco smoking - Never vs Former	6	944,931	OR 1.09 [1.01, 1.17]	I ² =90%	Very low
Tobacco smoking – Current or Former vs Never	12	985,086	OR 1.18 [1.10, 1.27]	I ² =81%	Very low

Abbreviations: CKD: chronic kidney disease; RR: relative risk; OR: odds ratio. ^a Figures in square brackets are 95% confidence intervals; ^b Summary of Findings table which details the reasons for downgrading evidence quality is reported in Supplemental Table 7.

Table 2. Meta-analyses performed in the review showing the association of alcohol consumption, diet, physical activity, and smoking exposure to secondary outcomes (kidney replacement therapy, GFR decline, and albuminuria)

Kidney Replacement Therapy^a					
Tobacco Smoking	Studies	Participants	Association ratio [95% CI]^b	Heterogeneity [I²]	Evidence Quality^c
Tobacco smoking - Former v current	7	1,126,913	RR 1.25 [1.13, 1.39]	I ² =39%	Moderate
Tobacco smoking - Current or Former vs Never	8	1,230,390	RR 1.59 [1.30, 1.94]	I ² =68%	Moderate
GFR decline^d					
Diet factors					
Potassium	4	10,729	RR 0.49 [0.31, 0.79]	I ² =81%	Very low
Protein	5	18,507	OR 1.07 [0.96, 1.19]	I ² =42%	Low
Physical Activity					
Physical activity - High vs Low levels	5	15,161	OR 0.77 [0.63, 0.93]	I ² =75%	Very low
Alcohol Consumption					
Alcohol - High vs Low intakes	5	16,580	OR 0.88 [0.72, 1.07]	I ² =15%	Very low
Albuminuria					
Diet Factors					
Sodium	3	11,751	OR 1.01 [0.89, 1.14]	I ² =0%	Very low
Physical Activity					
Physical activity - High vs Low levels	4	110,154	RR 0.88 [0.81, 0.96]	I ² =0%	Low
Alcohol consumption					
Alcohol - High vs Low intakes	7	220,479	RR 1.03 [0.88, 1.20]	I ² =59%	Low
Tobacco Smoking					
Tobacco smoking - Current or Former vs Never	7	184,302	OR 1.67 [1.23, 2.26]	I ² =88%	Very low

Abbreviations: GFR: glomerular filtration rate; RR: relative risk; OR: odds ratio ^a Meta-analysis not possible for diet factors, physical activity or alcohol consumption; ^b Figures in square brackets are 95% confidence intervals; ^c Summary of Findings table which details the reasons for downgrading evidence quality is reported in Supplemental Table 7; ^d Meta-analysis not possible for tobacco smoking.